Revisiting the Uranium-238 Thermal Neutron Capture Cross Section

A. Trkov¹, G.L. Molnár², Zs. Révay², S.F. Mughabghab³, R.B. Firestone⁴, V.G. Pronyaev¹, A.L. Nichols¹

Cross section databases have to be constantly improved in order to enhance further the predictive power of computational methods for reactor analysis and design. Clearly, the results are highly sensitive to the data adopted for the main uranium isotopes. One of the important recent changes introduced in the ENDF/B-VI library since Release 5 has been an increase of the capture to fission ratio of ²³⁵U in the epithermal energy range. This modification has improved the reactivity prediction in calculations for fast reactors, but caused under-predictions of reactivity in thermal lattices. Obviously, the previously low value for the epithermal capture of ²³⁵U compensated for an error in one or more other parameters, which still need to be identified. The thermal capture cross section of ²³⁸U is an obvious candidate for consideration. Within the Working Party on Evaluation Cooperation (WPEC), Subgroup 22 has been set up to address the problem. The function of the subgroup is primarily to co-ordinate activities and to exchange information, while the actual work is supported by national projects and other sources. Measurements of the thermal capture cross section of ²³⁸U are reviewed in the present work.

Several sets of ²³⁸U thermal capture cross section measurements from the literature were reviewed. Measurements by Hunt et al [HU69] and Poenitz et al [PO81] are consistent and require no corrections due to updated standards. The measurement by De Corte et al [DC88, DC92] is affected by the updated gamma-ray emission probability data; the corrected result is in very good agreement with the previous two studies. A new measurement by Molnár et al [MO02, MO03] was also analysed, which uses hydrogen as an internal standard. Although the uncertainty in this measurement is somewhat higher, the resulting data support the other measurements that are considered reliable. Table 1 summarises the results of the re-analysis.

The overall conclusion is that the Mughabghab [Mu84] recommendation for ²³⁸U thermal capture

Table 1: Summary of thermal capture cross sections of ²³⁸U after re-analysis.

Author	$oldsymbol{s}_{\!\gamma}[b]$	$\delta\sigma_{\gamma}$ [b]
Hunt et al [HU69]	2.69	0.03
Poenitz et al [PO81]	2.680	0.019
De Corte et al [DC88,DC92]	2.662	0.029
Molnár et al [MO02,MO03]	2.690	0.041
Adopted value	2.679	0.019

cross section of 2.680 ± 0.019 barns is valid. Old transmission measurements do not rely on neutron activation, and would not affect the average to a large extent because they are in good agreement with the present analysis. However, the recommended thermal cross section value for the ENDF/B-VI library was affected by the Bigham *et al* measurement [Bi69], which is too high. The measurement should be discarded and evaluated nuclear data files containing cross sections for ²³⁸U at thermal energies should be modified accordingly.

Footnotes and References

- ¹ Nuclear Data Section, IAEA
- ² Institute of Isotope and Surface Chemistry, Budapest
- ³ NNDC, Brookhaven National Laboratory
- ⁴ Lawrence Berkeley National Laboratory [HU69] J.B. Hunt, J.C. Robertson, T.B. Ryves, J. Nucl. Energy 23, (1969) 705-711.

[PO81] W.P. Poenitz, L.R. Fawcett, D.L. Smith, Nucl. Sci. Eng. 78, (1981) 239-347.

[DC88] F. De Corte, A. Simonits, J. Radioanal. Chem. 133, 1, (1989) 43.

[DC92] F. De Corte, J. Radioanal. Chem. 160, 1 (1992) 63-75.

[MO02] G.L. Molnar, Zs. Revay, T. Belgya, Nucl. Instrum. Meth. Phys. Res. B., in press.
[MO03] G.L. Molnar, private communication, 2003
[Bi69] Bigham et al, Can. J. Phys. 47, 1317 (1969)
[Mu84] S.F. Mughabghab, Neutron Cross Sections Vol. 1, Part B Z=61-100. (Academic Press, 1984).